THE ACTION FOR THREATENED MOTHS PROJECT

M. PARSONS, D. GREEN

Butterfly Conservation, UK Conservation Office, PO Box 444, Wareham, Dorset BH20 5YA.

P. WARING

1366 Lincoln Road, Werrington, Peterborough PE4 6LS.

HISTORICAL BACKGROUND

Butterfly Conservation (BC), a registered charity, was founded in 1968 as The British Butterfly Conservation Society. It now has a membership in excess of 8000 and is the largest invertebrate conservation society in Europe. Butterfly Conservation was set up with the aim of protecting our diminishing wild native butterflies and moths from destruction of habitat and other threats. The Society is lobbying continuously for a transformation in our attitudes to the countryside and its wildlife. The Society was entirely managed and run by volunteers until 1990 when an administration headquarters was set up. In 1993 the first full-time Conservation Officer was employed. The society now has over 15 staff and several contractors working on projects relating to Lepidoptera conservation.

Moths have always been included in the remit of BC, but until recently their profile was comparatively low within the Society. In 1993 BC contributed towards the National Moth Conservation Project, a tangible contribution that has continued to this day. The Society has contributed funds to several moth projects, including work on the bright wave *Idaea ochrata*, the marsh mallow *Hydraecia osseola*, the speckled footman *Coscinia cribraria* and the small dotted footman *Pelosia obtusa*, and funded an international conference on burnet moths. A number of regional workshops on moths have been organised for branches and there have been a wide range of moth related events held by the branches, e.g. several hundred moth trapping evenings. In 1999 the Society employed M. Parsons and D. Green on a full time basis with funding from English Nature (EN) and BC, along with P. Waring (PW) in an advisory role, to work on the Action for Threatened Moths Project and the National Recording Scheme for the Rarer British Macro-moths.

THE NATIONAL MOTH CONSERVATION PROJECT AND THE NATIONAL RECORDING SCHEME FOR THE RARER BRITISH MACROMOTHS

The National Moth Conservation Project was launched in 1987 by the former Nature Conservancy Council. This project has been operated by PW since its inception and includes work on species on Schedule 5 of the Wildlife & Countryside Act. There are several aims for this project. These include the formation and operation of a national information gathering network; to provide feedback to recorders; and to produce national distribution maps, particularly for the scarce and threatened species. This enables the regular assessment and revision of the conservation status of our scarce and threatened species. In 1990/91, the National Recording Scheme for the Rarer British Macro-moths was formed as part of the project by linking up existing County Moth Recorders and finding recorders for poorly covered areas. Further details and a resumé of the history of the project are given in Waring (1998 & 1999a). It is expected that the National Recording Scheme

BR. J. ENT. NAT H1ST., 13 2000

for the Rarer British Macro-moths will be integrated within the Action for Threatened Moths Project. This will provide a single point of contact and maximise the use of the data provided by contributors. Contributors will be informed of any change when it occurs. Ten annual news bulletins have been produced by the National Moth Conservation Project and sent to all County Moth Recorders. It is anticipated that future annual newsletters will be produced covering the National Moth Conservation Project, the National Recording Scheme for the Rarer British Macro-moths and the Action for Threatened Moths Project continuing PW's precedent.

BUTTERFLY CONSERVATION'S BRANCH MOTH OFFICERS AND THE COUNTY MOTH RECORDERS

Many of the current County Moth Recorders were individuals already working on county lists or providing some focal point for recording in a given county. The BC Branch Moth Officer post is relatively new. These started to be formally recognised and appointments made in 1994. *News bulletin 10* of the National Moth Conservation Project (Waring, 1999b & c) lists all the BC Branch Moth Officers and the County Moth Recorders. The role of the BC Branch Moth Officer is broadly to co-ordinate moth issues within the Branch and to promote moth recording and organise local events. This should involve forming and developing links with the County Moth Recorder and ensuring that all records of scarce or threatened moths from the Branch are forwarded to the National Recording Scheme for the Rarer British Macro-moths. Fuller details of the roles of BC Branch Moth Officers are given in Waring (1997). Branch Moth Officers and all other contributors are encouraged to send all records via the County Moth Recorder is also the Branch Moth Officer.

The UK biodiversity action plan and the action for threatened moths $$\operatorname{Project}$$

In response to the commitment given by the Prime Minister in signing the Convention on Biological Diversity at the Earth Summit in Rio de Janeiro in 1992. the UK Government published Biodiversity: The UK Action Plan (UK Biodiversity Group, 1994). In discharging our obligations under the Biodiversity Convention, the UK Action Plan set as an overall goal: "To conserve and enhance biological diversity within the UK and to contribute to the conservation of global biodiversity through all appropriate mechanisms". A Biodiversity Steering Group was established to advise government and to assist with work on biodiversity. In 1995 Volumes 1 and 2 of Biodiversity: The UK Steering Group Report (UK Biodiversity Group, 1995a & 1995b) were published. Volume 2 gave the Short, Middle and Long List of species being considered by the plan (this has since been amended) and included the Action Plans for three moths: the speckled footman Coscinia cribraria, netted carpet Eustroma reticulatum and the bright wave Idaea ochrata. Further Action Plans were published in UK Biodiversity Group (1999a,b). The preparation of all these documents drew heavily on the information collected by the National Moth Conservation Project.

There are a number of criteria, not all appropriate to moths, by which a species can qualify to be treated under the Biodiversity Action Plan (BAP). Although not comprehensive, the following criteria provide an indication of how the species were selected (after UK Biodiversity Group, 1995a).

- Threatened endemic and other globally threatened species;
- Species where the UK has more than 25% of the world or appropriate biogeographical population:
- Species where numbers or range have declined by more than 25% in the last 25 years;
- In some instances where the species is found in fewer than 15 10 km squares in the UK;
- Species which are listed in the EU Birds or Habitats Directives, the Bern, Bonn or CITES Conventions, or under the Wildlife and Countryside Act 1981 the Nature Conservation and Amenity Lands (Northern Ireland) Order 1985.

A summary of the Convention on Biological Diversity and how BC is working with a variety of partners within the framework of the convention is given by Bourn & Warren (1997).

The species in Table 1 are those covered by the BAP. In this species are covered by either a Priority Species Action Plan or a Species Statement. The Species Action Plans detail current status, current factors causing loss or decline, current action, objectives and targets and a range of proposed actions. The Species Statements are similar but under proposed action typically recommend monitoring only. For the purposes of the Action for Threatened Moths Project, the Species Action Plans and Statements are both treated equally. BC has agreed to be the Lead Partner for the projects on all but one of the priority moths and butterflies, sometimes in association with a statutory agency (e.g. EN) or a non-government organization (NGO) (e.g. the National Trust). It should be noted that several species covered by Schedule 5 of the Wildlife & Countryside Act are not included in this list. Work is expected to continue on these species under the Species Recovery Programme funded by EN or as projects of the Countryside Council for Wales (CCW) or Scottish Natural Heritage (SNH).

Prior to the publication of the majority of these Action Plans, BC started to produce a series of Regional Action Plans to identify regional priorities and draw together local information on priority butterflies and moths. Those already produced include Northern Ireland. Wales, West Midlands and North East England. These all include priorities for moth conservation in the regions, including many species that do not have a national Species Action Plan, and in many cases include details of ecology, distribution and actions necessary for their conservation. Implementing these plans is a major challenge for the Society, which will be aided by the Action for Threatened Moths Project.

The work undertaken within the Action for Threatened Moths Project is guided by a Steering Committee, currently made up with representatives of BC, the country agencies (i.e. EN, CCW, SNH and the JNCC) along with moth specialists. The main rationale behind the Project is to provide a co-ordinated approach to the conservation of the BAP moths and to increase the involvement of volunteers and other organisations in priority moth work. Initial work has included contacting all County Moth Recorders, county moth groups and BC Branch Moth Officers to inform them of the project. Some individuals have been contacted in order to encourage participation in preliminary survey work and new sites have already been discovered for several priority species.

THE FUTURE

The project aims to ensure that annual monitoring will be undertaken at key sites for most, if not all, of the BAP species. This may take the form of modified transects

† Acosmetia caliginosa reddish buff + (Hübn.) Aspitates gilvaria straw belle + (D. & S.) Athetis pallustris marsh moth + (Hübn.) Calophasia lunula toadflax brocade + (Hüfn.) Calophasia lunula toadflax brocade + (Hüfn.) dark crimson underwing + Catocala sponsa (L.) dark crimson underwing + Coleophora tricolor basil thyme case-bearer + Walsingham - - Coscinia cribraria (L.) speckled footman + Coscinia cribraria (L.) white-spotted pinion + Cyclophora pendularia dingy mocha + (Clerck) bart moth + Dicycla oo (L.) heart moth + Enstroma reticulatum netted carpet + (Borkhausen) - - Heliophobus reticulata bordered gothic + marginosa (Haw.) - - Hydraeia osseola marsh mallow + Hydraeia sylvata (D. & S.) waved carpet	Lead Partner
(D. & Š.)Athetis pallustrismarsh moth+(Hübn.)toadflax brocade+(Rüfn.)calophasia lunulatoadflax brocade+(Rüfn.)dark crimson underwing+Catocala sponsa (L.)dark crimson underwing+Catocala sponsa (L.)dark crimson underwing+Coscinia cribraria (L.)speckled footman+Coscinia cribraria (L.)speckled footman+Coscinia cribraria (L.)white-spotted pinion+Cyclophora pendulariadingy mocha+(Clerck)bart moth+Dicycla oo (L.)heart moth+Eustroma reticulatumnetted carpet+(D. & S.)nattwo bordered beauty+Hadena albinaculawhite spot+Heliophobus reticulatabordered gothic+marginosa (Haw.)narrow-bordered bee+Hydraeia osseolamarsh mallow+Hydraeia osseolamarsh mallow+Hydraeia osteolasilky wave+Hucherardi MabilleHypena rostralis (L.)buttoned snout+Lyceia zonaria britannicabelted beauty+Lyceia carbonariaclay fan-foot+Macaria carbonariaclay fan-foot+Minoa murinata (Scop.)scarce blackneck+Minoa murinata (Scop.)drab looper+Mona alpium (Osbeck)scarce mervielle du jour+Moria austaos (Hüb)lunar yellow under	BC
(Hübn.)Calophasia lunulatoadflax brocade(Hüfn.)Catocala promissa (D. & S.)light crimson underwingCatocala promissa (L.)dark crimson underwingCatocala promissa (L.)dark crimson underwingColeophora tricolorwalsinghamCoscinia cribraria (L.)speckled footmanCyclophora pendulariadingy mocha(Clerck)Dicycla oo (L.)heart mothpicycla oo (L.)heart moth(= paralellaria) (D. & S.)Eustroma reticulatumnetted carpet(Borkhausen)Heliophobus reticulatabordered gothicHadena albinaculaWaytaecia osseolamarginosa (Haw.)Henaris tityus (L.)narrow-bordered beeHydraecia osseolamarsh mallowHypena rostralis (L.)buttoned snoutHypena rostralis (L.)buttoned snoutHubhan,Idaea albinariasilky waveHubhan,Lypein corceago (D. & S.)scarce blacknecktypena rostralis (L.)buttoned snouttypena rostralis (L.)buttoned snoutthypena rostralis (L.)	BC
Calophasia hundatoadflax brocade+(Hüfn.)toadflax brocade+Catocala promissa (D. & S.)light crimson underwing+Catocala sponsa (L.)dark crimson underwing+Catocala sponsa (L.)dark crimson underwing+Coleophora tricolorbasil thyme case-bearer+WalsinghamCoscinia cribraria (L.)speckled footman+Cosmia diffinis (L.)white-spotted pinion+Cophora pendulariadingy mocha+(Clerck)Dicycla oo (L.)heart moth+(= paralellaria) (D. & S.)Eustroma reticulatumnetted carpet+(Borkhausen)Hadema albinaculawhite spot+Heliophobus reticulatabordered gothic+marginosa (Haw.)Hydrelia sylvata (D. & S.)waved carpet+Hydraecia osseolaHydena rostralis (L.)buttoned snout+Idaea oclirata (Scop.)orange upperwing+Lycia zonaria britannicabelted beauty+(Hübn.)Idaea oclirata (Scop.)scarce marvielle du jour+Mona alpium (Osbeck)scarce mervielle du jour+Mona alpium (Osbeck) <td>BC</td>	BC
Catocala promissa (D. & S.)light crimson underwing+Catocala sponsa (L.)dark crimson underwing+Catocala sponsa (L.)basil thyme case-bearer+Coleophora tricolorbasil thyme case-bearer+WalsinghamCoscinia cribraria (L.)speckled footman+Coscinia cribraria (L.)white-spotted pinion+Coscinia diffinis (L.)white-spotted pinion+Corck)Dicycla oo (L.)heart moth+Environe vespertaria (L.)dark bordered beauty+(- paradellaria) (D. & S.)Eustrona reticulatumnetted carpet+(Borkhausen)Heliophobus reticulatabordered gothic+Hydraetia sylvata (D. & S.)Hydraetia sylvata (D. & S.)waved carpet+Hydraetia sylvata (D. & S.)Hydraetia sylvata (D. & S.)buttoned snout+Hydraetia sylvata (D. & S.)Hucherardi MabilleHypena rostralis (L.)buttoned snout+Idaea ochrata (Scop.)bright wave+Idaea ochrata (Scop.)bright wave+Idaea ochrata (Scop.)Hacania carbonariaLycic zonaria britannicabelted beauty+Lycic zonaria britannicabelted beauty+Hydraecia carbonariaClerck) </td <td>BC</td>	BC
Catocala sponsa (L.)dark crimson underwing+Coleophora tricolorbasil thyme case-bearer+Walsingham-Coscinia cribraria (L.)speckled footman+Coscinia cribraria (L.)white-spotted pinion+Coyclophora pendulariadingy mocha+(Clerck)-dirk bordered beauty+Dicycla oo (L.)heart moth+Eustrona reticulationnetted carpet+(D. & S.)+Hadena albinaculawhite spot+(Borkhausen)Heliophobus reticulatabordered gothic+Hydrelia sylvata (D. & S.)Hydraecia osseolamarsh mallow+Hydraecia osseolaHypena rostralis (L.)buttoned snout+Jodia croceago (D. & S.)Idaea albinanicabelted beauty+(Hübn.)Idaea alchata (Scop.)bright wave+Jodia croceago (D. & S.)scarce blackneck+Macaria carbonarianetted mountain moth+Lycia zonaria britannicabelted beauty+(Clerck)Minoa numinata (Scop.)drab looper+Moma alpinm (Osbeck)scarce mervielle du jour+Moma alpinm (Osbeck)scarce mervielle du jour+Paracolax tristalis (Fab.)clay fan-foot+	BC
Coleophora tricolorbasil thyme case-bearer+Walsingham-Coscinia cribraria (L.)speckled footman+Cosnia diffinis (L.)white-spotted pinion+Cyclophora pendulariadingy mocha+(Clerck)heart moth+Dicycla oo (L.)heart moth+Epione vespertaria (L.)dark bordered beauty+(Clerck)netted carpet+(D. & S.)reticulatumnetted carpet+(Borkhausen)bordered gothic+Heliophobus reticulatabordered gothic+Marginosa (Haw.)narrow-bordered bee+Hydraecia osseolamarsh mallow+Hydraecia osseolamarsh mallow+Hypena rostralis (L.)buttoned snout+Idaea dilutariasilky wave+(Hübn.)bright wave+Idaea ochrata (Scop.)bright wave+(Clerck)Macaria carbonaria-+Macaria carbonaria-+Macaria carbonaria-+Macaria carbonaria-+Macaria carbonaria+-Mana alpium (Osbeck)scarce mervielle du jour+Mona alpium (Osbeck)scarce mervielle du jour+Mortia arbona Hüfn.)lunar yellow underwing+Mortia arbona Hüfn.)lunar yellow underwing+Paracolax tristalis (Fab.)clay fan-foot+	BC
Coscinia cribraria (L.)speckled footman+Cosmia diffinis (L.)white-spotted pinion+Cosmia diffinis (L.)white-spotted pinion+Cyclophora pendulariadingy mocha+(Clerck)beart moth+Dicycla oo (L.)heart moth+(Elerck)dark bordered beauty+(= paralellaria) (D. & S.)beart moth+Eustroma reticulatumnetted carpet+(D. & S.)heart moth+Hadena albimaculawhite spot+(Borkhausen)bordered gothic+Heliophobus reticulatabordered gothic+hawk-mothhawk-moth+Hydraetia sylvata (D. & S.)marrow-bordered bee+Hydraetia sylvata (D. & S.)waved carpet+Hydraetia sylvata (D. & S.)buttoned snout+Hydraetia sylvata (Scop.)bright wave+Idaea ochrata (Scop.)bright wave+Jodia croceago (D. & S.)orange upperwing+Lyceia zonaria britannicabelted beauty+Lygephila craccae (D. & S.)scarce blackneck+Macaria carbonarianetted mountain moth+(Clerck)drab looper+Minoa murinata (Scop.)drab looper+Mona alpium (Osbeck)scarce mervielle du jour+Moria alpium (Osbeck)scarce mervielle du jour+Paracolax tristalis (Fab.)clay fan-foot+	BC
Cosmia diffinis (L.)white-spotted pinion+Cyclophora pendulariadingy mocha+Cyclophora pendulariadingy mocha+Clerck)bicycla oo (L.)heart moth+Dicycla oo (L.)heart moth+Epione vespertaria (L.)dark bordered beauty+(= paraleilaria) (D. & S.)netted carpet+Eustroma reticulatumnetted carpet+(D. & S.)narrow-bordered bee+Hadena albinaculawhite spot+(Borkhausen)bordered gothic+Heliophobus reticulatabordered gothic+hawk-mothhawk-mothHydraeia sylvata (D. & S.)waved carpet+Hydraeia osseolamarsh mallow+luncherardi Mabille+-Hypena rostralis (L.)buttoned snout+Idaea dilutariasilky wave+Jodia croccago (D. & S.)orange upperwing+Lycia zonaria britannicabelted beauty+(Harrison)Lygephila craccae (D. & S.)scarce blackneck+Minoa nurinata (Scop.)drab looper+Minoa nurinata (Scop.)drab looper+Moma alpium (Osbeck)scarce mervielle du jour+Mortua orbona Hüfn.)lunar yellow underwing+Noctua orbona Hüfn.)lunar yellow underwing+Paracolax tristalis (Fab.)clay fan-foot+	BC
Cyclophora pendulariadingy mocha+(Clerck)heart moth+Dicycla oo (L.)heart moth+Epione vespertaria (L.)dark bordered beauty+(= paralellaria) (D. & S.)netted carpet+(Estroma reticulatumnetted carpet+(D. & S.)natted carpet+(Borkhausen)bordered gothic+Heliophobus reticulatabordered gothic+Maginosa (Haw.)narrow-bordered bee+Hemaris tityus (L.)narrow-bordered bee+Hydrelia sylvata (D. & S.)waved carpet+Hydraecia osseolamarsh mallow+Hypena rostralis (L.)buttoned snout+Idaea olintariasilky wave+(Hübn.)ldaea ochrata (Scop.)bright waveLycia zonaria bitannicabelted beauty+Lygephila craceae (D. & S.)scarce blackneck+Macaria carbonarianetted mountain moth+(Clerck)drab looper+Mona alpinm (Osbeck)scarce mervielle du jour+Mythinna turca (L.)double line+Noctua orbona Hüfn.)lunar yellow underwing+Oria musculosa (Hübn)brighton wainscot+	BC
Epione vespertaria (L.)dark bordered beauty+(= paralellaria) (D. & S.)netted carpet+(D. & S.)netted carpet+(D. & S.)Hadena albinaculawhite spot+Hadena albinaculawhite spot+(Borkhausen)bordered gothic+Heliophobus reticulatabordered gothic+marginosa (Haw.)narrow-bordered bee+Hemaris tityus (L.)narrow-bordered bee+Hydraecia osseolamarsh mallow+Hucherardi Mabille+Hypena rostralis (L.)buttoned snout+Idaea dilutariasilky wave+Jodia croceago (D. & S.)orange upperwing+Lycia zonaria britannicabelted beauty+Lycai zonaria britannicabelted beauty+Macaria carbonarianetted mountain moth+(Clerck)drab looper+Mona alpium (Osbeck)scarce mervielle du jour+Mona alpium (Osbeck)scarce mervielle du jour+Mona alpium (Osbeck)scarce mervielle du jour+Mona alpium (Sbeck)scarce mervielle du jour+Paracolax tristalis (Fab.)clay fan-foot+	BC
(= paralellaria) (D. & S.)Eustroma reticulatumnetted carpet(D. & S.)Hadena albimaculawhite spotHadena albimaculawhite spot(Borkhausen)Heliophobus reticulatabordered gothic+marginosa (Haw.)Hemaris tityus (L.)narrow-bordered bee+hukemothHydraecia osseolamarsh mallowHypena rostralis (L.)buttoned snoutHalaea dilutariasilky wave(Hübn.)Idaea ochrata (Scop.)bright waveHarrison)Lycia zonaria britannicabelted beauty+Macaria carbonarianetted mountain moth+Mona alpium (Osbeck)scarce mervielle du jour+Mythinna turca (L.)double line+Oria nusculosa (Hübn)brighton wainscot+Paracolax tristalis (Fab.)clay fan-foot+	BC
Enstroma reticulatumnetted carpet+(D. & S.)white spot+(Borkhausen)bordered gothic+Heliophobus reticulatabordered gothic+marginosa (Haw.)narrow-bordered bee+Hemaris tityus (L.)narrow-bordered bee+hawk-mothhawk-mothHydrelia sylvata (D. & S.)waved carpet+Hydraecia osseolamarsh mallow+hucherardi Mabille+Hypena rostralis (L.)buttoned snout+Idaea dilutariasilky wave+Idaea ochrata (Scop.)bright wave+Jolia croceago (D. & S.)orange upperwing+Lygephila craccae (D. & S.)scarce blackneck+Macaria carbonarianetted mountain moth+(Clerck)drab looper+Mona alpinm (Osbeck)scarce mervielle du jour+Monia alpinm (Osbeck)scarce mervielle du jour+Mortua orbona Hüfn.)lunar yellow underwing+Oria musculosa (Hübn)brighton wainscot+Paracolax tristalis (Fab.)clay fan-foot+	BC/ RSPB
Hadena albimaculawhite spot+(Borkhausen)bordered gothic+Heliophobus reticulatabordered gothic+marginosa (Haw.)narrow-bordered bee+Hemaris tityus (L.)narrow-bordered bee+hawk-mothhawk-mothHydrelia sylvata (D. & S.)waved carpet+Hydraecia osseolamarsh mallow+hucherardi Mabille+Hypena rostralis (L.)buttoned snout+Idaea dilutariasilky wave+Idaea ochrata (Scop.)bright wave+Jodia croceago (D. & S.)orange upperwing+Lycia zonaria britannicabelted beauty+(Harrison)Lygephila craccae (D. & S.)scarce blackneck+Mona alpinm (Osbeck)scarce mervielle du jour+Moma alpinm (Osbeck)scarce mervielle du jour+Monta alpinm (Usbeck)scarce mervielle du jour+Paracolax tristalis (Fab.)clay fan-foot+	BC/NT
Heliophobus reticulata marginosa (Haw.)bordered gothic+Hemaris tityus (L.) hawk-mothnarrow-bordered bee hawk-mothHydrelia sylvata (D. & S.)waved carpetHydraecia osseola hucherardi Mabillemarsh mallowHypena rostralis (L.)buttoned snoutHaea dilutaria occeago (D. & S.)silky waveHaea ochrata (Scop.)bright waveHycia croceago (D. & S.) orange upperwing+Lycia zonaria britannica belted beauty+Lygephila craccae (D. & S.) (Clerck)scarce blackneckMinoa murinata (Scop.)drab looperMinoa alpium (Osbeck) 	BC
Hemaris tityus (L.)narrow-bordered bee hawk-mothHydrelia sylvata (D. & S.)waved carpetHydraecia osseolamarsh mallowhucherardi MabilleHypena rostralis (L.)buttoned snoutHaea dilutariasilky wave(Hübn.)Idaea ochrata (Scop.)bright waveIdaea ochrata (Scop.)bright waveLycia zonaria britannicabelted beautyLygephila craccae (D. & S.)scarce blackneckMacaria carbonarianetted mountain moth(Clerck)tab looperMinoa murinata (Scop.)drab looperMonua alpinm (Osbeck)scarce mervielle du jour+double line+houle line+found artifich.)Iunar yellow underwing+priaton wainscot+priaton wainscot+hughton wainscot++hugh	BC
Hydrelia sylvata (D. & S.)waved carpet+Hydraecia osseolamarsh mallow+hucherardi Mabillemarsh mallow+Hypena rostralis (L.)buttoned snout+Idaea dilutariasilky wave+(Hübn.)bright wave+Idaea ochrata (Scop.)bright wave+Jodia croceago (D. & S.)orange upperwing+Lycia zonaria britannicabelted beauty+(Harrison)Lygephila craccae (D. & S.)scarce blackneck+Macaria carbonarianetted mountain moth+(Clerck)Minoa murinata (Scop.)drab looper+Monta alpinm (Osbeck)scarce mervielle du jour+Noctua orbona Hüfn.)lunar yellow underwing+Oria musculosa (Hübn)brighton wainscot+Paracolax tristalis (Fab.)clay fan-foot+	BC
Hydraecia osseola hucherardi Mabillemarsh mallow+Hydraecia osseola hucherardi Mabillemarsh mallow+Hypena rostralis (L.)buttoned snout+Idaea dilutaria silky wavesilky wave+Idaea dilutaria (Hübn.)silky wave+Idaea ochrata (Scop.)bright wave+Jodia croceago (D. & S.) (Harrison)orange upperwing+Lycia zonaria britannica (Harrison)belted beauty+Lygephila craccae (D. & S.) (Clerck)scarce blackneck+Macaria carbonaria (Clerck)netted mountain moth+Mona alpium (Osbeck) Noctua orbona Hüfn.)scarce mervielle du jour+Mythinna turca (L.) Oria nusculosa (Hübn)brighton wainscot+Paracolax tristalis (Fab.)clay fan-foot+	BC
Idaea dilutaria silky wave + (Hübn.) bright wave + Idaea ochrata (Scop.) bright wave + Jodia croceago (D. & S.) orange upperwing + Lycia zonaria britannica belted beauty + (Harrison) - - Lygephila craccae (D. & S.) scarce blackneck + Macaria carbonaria netted mountain moth + (Clerck) - + Minoa murinata (Scop.) drab looper + Mona alpium (Osbeck) scarce mervielle du jour + Mythinna turca (L.) double line + Oria musculosa (Hübn) brighton wainscot + Paracolax tristalis (Fab.) clay fan-foot +	BC
Idaea dilutariasilky wave+(Hübn.)Idaea ochrata (Scop.)bright wave+Idaea ochrata (Scop.)bright wave+Jodia croceago (D. & S.)orange upperwing+Lycia zonaria britannicabelted beauty+(Harrison)Lygephila craceae (D. & S.)scarce blackneck+Macaria carbonarianetted mountain moth+(Clerck)Minoa murinata (Scop.)drab looper+Monna alpium (Osbeck)scarce mervielle du jour+Nottua orbona Hüfn.)lunar yellow underwing+Oria musculosa (Hübn)brighton wainscot+Paracolax tristalis (Fab.)clay fan-foot+	BC
Jodia croceago (D. & S.)orange upperwing+Lycia zonaria britannicabelted beauty+(Harrison)Lygephila craceae (D. & S.)scarce blackneck+Macaria carbonarianetted mountain moth+(Clerck)Minoa murinata (Scop.)drab looper+Moma alpinm (Osbeck)scarce mervielle du jour+Mythinna turca (L.)double line+Noctua orbona Hüfn.)lunar yellow underwing+Oria musculosa (Hübn)brighton wainscot+Paracolax tristalis (Fab.)clay fan-foot+	BC
Jodia croceago (D. & S.) orange upperwing + Lycia zonaria britannica belted beauty + (Harrison) + + Lygephila craceae (D. & S.) scarce blackneck + Macaria carbonaria netted mountain moth + (Clerck) - - Minoa murinata (Scop.) drab looper + Mythimna turca (L.) double line + Noctua orbona Hüfn.) lunar yellow underwing + Oria musculosa (Hübn) brighton wainscot + Paracolax tristalis (Fab.) clay fan-foot +	BC
Lycia zonaria britannica belted beauty + (Harrison) telted beauty + Lygephila craccae (D. & S.) scarce blackneck + Macaria carbonaria netted mountain moth + (Clerck) - - Minoa murinata (Scop.) drab looper + Moma alpium (Osbeck) scarce mervielle du jour + Mythinna turca (L.) double line + Noctua orbona Hüfn.) lunar yellow underwing + Oria musculosa (Hübn) brighton wainscot + Paracolax tristalis (Fab.) clay fan-foot +	BC
Macaria carbonaria (Clerck) netted mountain moth + Minoa nurinata (Scop.) drab looper + Mona alpium (Osbeck) scarce mervielle du jour + Mythinna turca (L.) double line + Noctua orbona Hüfn.) lunar yellow underwing + Oria nusculosa (Hübn) brighton wainscot + Paracolax tristalis (Fab.) clay fan-foot +	BC
(Clerck)drab looper+Minoa murinata (Scop.)drab looper+Moma alpium (Osbeck)scarce mervielle du jour+Mythinma turca (L.)double line+Noctua orbona Hüfn.)lunar yellow underwing+Oria musculosa (Hübn)brighton wainscot+Paracolax tristalis (Fab.)clay fan-foot+	BC
Minoa murinata (Scop.)drab looper+Moma alpium (Osbeck)scarce mervielle du jour+Mythimma turca (L.)double line+Noctua orbona Hüfn.)lunar yellow underwing+Oria musculosa (Hübn)brighton wainscot+Paracolax tristalis (Fab.)clay fan-foot+	BC
Moma alpium (Osbeck)scarce mervielle du jour+Mythimma turca (L.)double line+Noctua orbona Hüfn.)lunar yellow underwing+Oria musculosa (Hübn)brighton wainscot+Paracolax tristalis (Fab.)clay fan-foot+	BC
Noctua orbona Hüfn.)lunar yellow underwing+Oria musculosa (Hübn)brighton wainscot+Paracolax tristalis (Fab.)clay fan-foot+	BC
Oria musculosa (Hübn)brighton wainscot+Paracolax tristalis (Fab.)clay fan-foot+	BC
Paracolax tristalis (Fab.) clay fan-foot +	BC
	BC
(= derivalis (Hübn.))	BC
Paradiarisa sobrina (Dup.) cousin german +	BC

Table 1: Species covered by the Biodiversity Action Plan

(Continued)

BR. J. ENT. NAT. HIST., 13: 2000

Table 1. (Continued)

Species	English Name	Priority Species Action Plan	Species Statement	Lead Partner
<i>†Pareulype berberata</i> (D. & S.)	barberry carpet	+		BC
Pechipogo strigilata (L.)	common fan-foot	+		BC
Phyllodesma ilicifolia (L.)	small lappet		+	BC
Polia bombycina (Hufn.)	pale shining brown	+		BC
Polymixis xanthomista (Hübn.)	black-banded		+	BC
<i>Pyropteron chrysidiformis</i> (Esper)	fiery clearwing	+		BC/EN
Rheumaptera hastata (L.)	argent & sable	+		BC
Schrankia taenialis (Hübn.)	white-line snout		+	BC
Scotopteryx bipunctaria (D. & S.)	chalk carpet		+	BC
Shargacucullia lychnitis (Rambur)	striped lychnis	+		BC
Siona lineata (Scop.)	black-veined moth	+		BC
<i>†Thetidia smaragdaria</i> <i>maritima</i> (Prout)	essex emerald		+	BC
Trichopteryx polycommata (D. & S.)	barred tooth-striped	+		BC
Trisateles emortualis (D. & S.)	olive crescent		+	BC
Tyta huctuosa (D. & S.)	four-spotted		+	BC
Xestia alpicola alpina (Humphreys & Westwood)	northern dart		+	BC
Xestia ashworthii (Doubleday)	ashworth's rustic		+	BC
Xestia rhomboidea (Esper)	square-spotted clay	+		BC
Xylena exsoleta (L.)	sword-grass	+		BC
Zygaena loti scotica (Rowland-Brown)	slender scotch burnet	+		BC
†Zygaena viciae argyllensis Tremewan	new forest burnet	+		SNH

Key: BC Butterfly Conservation; EN English Nature; NT National Trust; RSPB Royal Society for the Protection of Birds; SNH Scottish Natural Heritage; †Wildlife and Countryside Act species

after Pollard (1977), Spalding (1997) and Birkinshaw & Thomas (1999), thorough light trapping or by larval surveys. This monitoring, however, will depend primarily on volunteer effort and future funding for specific work. Further surveys will also take place aimed at understanding the distribution of individual species and identifying priorities for further effort. In some cases extensive autecological research may be needed and substantial additional funds will be required. Fund-raising is thus a vital part of our role in conjunction with specialist staff within BC.

Practical conservation measures are expected to be implemented for many of the BAP species, though in some cases our understanding of an individual species' ecology will need to improve before any measures can be put in place. Part of the

process will be to ensure that the local teams of the conservation agencies and land owners are aware of the presence of BAP species.

As is inherent in the BAP process, review of the current list of species will be ongoing. If it becomes obvious that other species have been overlooked and are in need of conservation effort, particularly if they meet the aforementioned criteria, an Action Plan will be drafted and effort made for it to be implemented. With the increase in interest and the publication of a number of key works, such as The Moths and Butterflies of Great Britain and Ireland series, the knowledge of the so-called micro-lepidoptera is such that conservation of some may benefit from the formulation and implementation of Action Plans. With an active scheme such as the Pyralid & Plume Recording Scheme run by Tony Davis and the existence of a national review (Parsons 1993), it is comparatively straightforward to identify several species of pyralid moth that are candidates for this approach. It may not be quite so straightforward to identify other candidate species, but if other species reviews are undertaken this task should get easier. In the meantime certain species suggest themselves through habitat concerns, e.g. Coleophora vibicella on dyer's greenweed Genista tinctoria and C. ochrea on common rock-rose Helianthemum nummularium growing in exposed situations.

The Council of BC has recently revised the Society's Reserve Acquisition Strategy. Potential reserves will have to meet various criteria, including safeguarding the wildlife interest (for example, land becoming available and threatened through a change in management). education opportunities and location. As opportunities arise, reserves may be established and purchased primarily for their moth interest.

It is expected that increasing use will be made of computer technology, but this will not exclude any individuals from contributing to any aspect of the project. For example, we have been trialing a weekly newsletter sent out via e:mail to surveyors aimed particularly at increasing the recording of the BAP species. It is anticipated that there will be an increased use of e:mail for correspondence (our e:mail addresses are mparsons@butterfly-conservation.org and dgreen@butterfly-conservation.org).

HOW YOU CAN HELP

Various surveys and monitoring projects are planned and we need volunteers to undertake aspects of these. In the future there may be also the opportunity to become involved with some more intensive autecological work. In the meantime, recording under-worked areas can result in unexpected finds, which can add considerably to our knowledge of individual species. Please ensure that all records are forwarded to the County Moth Recorder, from whom all relevant records should be forwarded to us for future incorporation into the National Recording Scheme for the Rarer British Macro-moths. We hope that surveys for BAP species can be organised and promoted at a local level, for example by the BC Branch Moth Officer or the County Moth Recorder. Obviously, the more people that participate in these surveys then the more comprehensive the results.

It is hoped that this process will also bring Societies such as BC and the British Entomological & Natural History Society and the local BC Branches and the various moth or invertebrate groups into much closer contact and, perhaps, co-operation with individual projects. This will undoubtedly have benefits for all concerned and for moth conservation.

If you have any comments on any aspect of this article or the project we would be pleased to hear from you.

ACKNOWLEDGEMENTS

We would like to take the opportunity to thank Martin Warren, Nigel Bourn, David Bridges and Roger Smith for useful comments on earlier drafts of the text.

References

- Birkinshaw, N. & Thomas, C.D. 1999. Torch-light transect surveys for moths. Journal of Insect Conservation, 3: 15–24.
- Bourn, N. & Warren, M. 1997. The road from Rio: Conservation action for the UK's priority butterflies. *Butterfly Conservation News*, No. 65: 7–9.
- Parsons, M.S. 1993. A review of the scarce and threatened pyralid moths of Great Britain. Joint Nature Conservation Committee, Peterborough. (UK Nature Conservation No. 11).
- Pollard, E. 1977. A method for assessing changes in the abundance of butterflies. *Biological Conservation*, 12: 115-134.
- Spalding, A. 1997. The use of the butterfly transect method for the study of the nocturnal moth Luperina nickerlii leechi Goater (Lepidoptera: Noctuidae) and its possible application to other species. Biological Conservation, 80: 147–152.
- UK Biodiversity Group, 1994. Biodiversity. The UK Action Plan. HMSO, London.
- UK Biodiversity Group, 1995a. Biodiversity: The UK Steering Group Report. Volume 1: Meeting the Rio Challenge. HMSO, London.
- UK Biodiversity Group, 1995b. Biodiversity: The UK Steering Group Report. Volume 2: Action Plans. HMSO, London.
- UK Biodiversity Group, 1999a. UK Biodiversity Group. Tranche 2 Action Plans. Volume IVinvertebrates. English Nature, Peterborough.
- UK Biodiversity Group, 1999b. UK Biodiversity Group. Tranche 2 Action Plans. Volume VIterrestrial and freshwater species and habitats. English Nature, Peterborough.
- Waring, P. 1997. The main roles of the Butterfly Conservation Branch Moth Officers. National Moth Conservation Project. News bulletin, 8: 7–8.
- Waring, P. 1998. A resume of the history, operation and products of the work of the National Recording Network for the Rarer British Macro-Moths and ideas on developments for the future. National Moth Conservation Project. *News bulletin*, 9: 7–12.
- Waring, P. 1999a. The National Recording Network for the Rarer British Macro-moths. Atropos, 6: 19-27.
- Waring, P. 1999b. Directory to the names and addresses of the County Moth Recorders. National Moth Conservation Project. News bulletin, 10: [33]-[36].
- Waring, P. 1999c. An up-dated directory to the names and addresses of Butterfly Conservation's Branch Moth Officers. National Moth Conservation Project. *News bulletin*, 10: [36]–[38].

BUTTERFLY CONSERVATION—ACTION FOR THREATENED MOTHS PROJECT

AN OPPORTUNITY FOR PARTICIPATION BY BENHS MEMBERS

THE BUTTERFLY CONSERVATION PROJECT

Butterfly Conservation (BC) are undertaking a major new project which will concentrate conservation action on the UK's most threatened moth species. This project is in part grant-aided by English Nature (EN).